

NICU Vascular Access Guidelines

Background:

Vascular access in neonates is frequently required to provide IV fluids, medications, and blood products. Determining the proper type of access for each patient and using the access site appropriately is critical in providing safe and effective care.

Indication:

Common NICU Vascular Access Options

	UVC	UAC	PICC	PAL	PIV
Indications:	1. Difficult PIV access 2. Hemodynamic instability 3. High GIR need	1. Hemodynamic instability 2. Frequent blood draws 3. Maintaining Arterial Access for future Cath Procedures in infants with Congenital Heart Disease (CHD)	1. Needed for IVF or medications (Unable to place UVL or UVL no longer useable)	1. Hemodynamic instability 2. Frequent blood draws (Unable to place UAL)	1. Needed for IVF or medications 2. Needed for blood products
Gestational Age Considerations					

<26 weeks	Recommended	Recommended	Consider if indicated (see above)	Consider if indicated (see above)	Consider if indicated (see above)
27-30 weeks	Recommended	Consider if indicated (see above)	Consider if indicated (see above)	Consider if indicated (see above)	Consider if indicated (see above)
>30 weeks	Consider if indicated (see above)	Consider if indicated (see above)	Consider if indicated (see above)	Consider if indicated (see above)	Consider if indicated (see above)



Placement/ Position:

Type: Umbilical Venous Catheter (UVC)

UVC	
Size	< 1.5kg: 3.5F >1.5kg: 5F
Length	$[(\text{Wt. in kg} \times 3 + 9) \div 2] + 1 + \text{length of stump (cm)}$
Placement	Target: ~T8-T9 (Junction of IVC and Right Atrium) Avoid: Right Atrium (above T7), Portal Veins (~T11) Low Lying UVC (use should be avoided when possible): Tip terminates below liver
Confirmation of Position	<u>*Initial:</u> Babygram + Cross-table <u>*Follow-Up:</u> Babygram 6-12 hours later <u>*After Adjustments:</u> If line moved $\geq 0.5\text{cm}$, repeat Xray within 30-60 min

	Image obtained from: https://torontocentreforneonatalhealth.com/wp-content/uploads/2019/07/Umbilical-Catheters-and-Peripherally-Inserted-Central-Catheters-PICC-Guideline.pdf
May use for	IVF: D10 (+/- electrolytes), TPN/IL, NS boluses, Must + Add Heparin 0.5 units/ml Meds: All meds ok Blood Products: All Blood products ok TKO Fluid: Must + Add Heparin 0.5 units/ml [For Low Lying UVLs: Only use non-hyperosmolar fluids, attending approval for anything else is required]
Duration of use	* Remove as soon as no longer needed * May be used up to 7 days * If line is determined critical, needs attending approval for 7+ days * For ALL infants with Congenital Heart Disease (CHD), discuss need for continued UVC use with Cardiology/Interventional Cardiology PRIOR to removing line


Type: Umbilical Arterial Catheter (UAC)

UAC	
Size	< 1.5kg: 2.5 Fr, 3.5F >1.5kg: 5F
Length	(Wt in kg x 3 +9) + length of stump (cm)
Placement	Target: ~T6-T9 Acceptable: L3-4 Avoid: T10-L2, Below L4 (Bifurcation of the Aorta)
Confirmation of Position	<u>*Initial:</u> Babygram <u>*Follow-Up:</u> Babygram 6-12 hours later <u>*After Adjustments:</u> If line moved $\geq 0.5\text{cm}$, repeat Xray within 30-60 min   Image obtained from: https://torontocentreforneonatalhealth.com/wp-content/uploads/2019/07/Umbilical-Catheters-and-Peripherally-Inserted-Central-Catheters-PICC-Guideline.pdf
May use for	IVF: Must + Add Heparin 0.5units/ml Meds: None

	Blood Products: Not routinely used for blood administration TKO Fluid: Must + Add Heparin 0.5units/ml
Duration of use	* Remove as soon as no longer needed or if any clinical signs of vascular insufficiency * May be used up to 5-7 days * If line is determined critical, needs attending approval for 7+ days * For ALL infants with Congenital Heart Disease (CHD), discuss need for continued UAC use with Cardiology/Interventional Cardiology PRIOR to removing line

Type: Peripheral Inserted Central Catheter (PICC)

PICC	
Size	Dependent on vessel size Neonatal PICC: Typically ≤ 1.9 Fr (1.4Fr Single Lumen, 1.9Fr Single Lumen, 1.9Fr Double Lumen) Pediatric PICC: For larger infants, attendings can request Pediatric PICC team to assess an infant's eligibility for a larger catheter central line.
Length	Catheter trimmed to size by person inserting it
Placement	<u>Right Upper Extremity:</u> Target: T4-5 Avoid: Right Atrium (~below T5)



Left Upper Extremity:

Target: T4-5

Avoid: Right Atrium (~below T5)





Image obtained from: <https://torontocentreforneonatalhealth.com/wp-content/uploads/2019/07/Umbilical-Catheters-and-Peripherally-Inserted-Central-Catheters-PICC-Guideline.pdf>

Lower Extremity:

Target: T 9-11

Acceptable: Above L2

	Avoid: Right Atrium T7, Renal Veins L2, Below L3-4 (Bifurcation of the Aorta)  Image obtained from: https://torontocentreforneonatalhealth.com/wp-content/uploads/2019/07/Umbilical-Catheters-and-Peripherally-Inserted-Central-Catheters-PICC-Guideline.pdf
Confirmation of Position	<u>Initial:</u> Upper Extremity: Chest X-ray, both shoulders abducted 30* Lower Extremity: Babygram x-ray, frog leg position <u>Follow-Up:</u> 6-12 hours later <u>After Adjustments:</u> If line moved ≥ 0.5cm, repeat Xray within 30-60 min <u>Maintenance:</u> Weekly Monday Xrays for picc line placement confirmation <u>Transduce:</u> Bedside RN to transduce PICC line to ensure it's not arterial before use
May use for	IVF: Must + Add Heparin 0.5units/ml Meds: All Blood Products: None TKO Fluid: Run at 1ml/hr. (May be run at 0.8ml/hr or 0.5ml/hr with attending approval)
Duration of use	* Remove as soon as no longer needed

Ellucid Policy	See Policy “Vascular Access Policy (Neonatal) 13002” for additional PICC line resources
----------------	---

Type: Peripheral Arterial Line

PAL	
Size	24G Angiocath
Length	n/a
Placement	Radial Artery (Left is pre-ductal, Right is post-ductal) Dorsal Pedis Artery Posterior Tibial Artery
Confirmation of Position	Ability to flush and draw blood easily Arterial waveform on Monitor
May use for	IVF: No Meds: No Blood Products: No TKO Fluid: 0.5ml/hr of Sterile Water + Na Acetate + Lidocaine + Heparin
Duration of use	* Remove as soon as no longer needed

Other types of lines used in the NICU:

Type	Notes
Pediatric PICC:	- Placed by Pediatric PICC Team - Size: 3Fr Single Lumen, 4Fr Double Lumen - May be used for Blood draws and Blood product transfusions
Broviac:	- Placed by Pediatric Surgery - Size: 2.7Fr Single Lumen - Can be used for Blood draws and Blood product transfusions
Femoral or Internal Jugular (IJ) Central Line:	- Placed by PICU attending or Ped Surgery - Size: 3Fr Single Lumen, 4Fr Double Lume - Can be used for Blood draws and Blood product transfusions

Infection Prevention:

Dressing care:

- See Ellucid Policy 13002: "Vascular Access Policy (Neonatal)" for additional PICC dressing Change instructions.
- Dressing should be intact, clean, and dry.



- To minimize the risk of line migration and infection, PICC dressings must be changed as soon as loss of dressing integrity is noted.
 - o a) For infants 36 weeks corrected gestational age and greater without a CHG allergy, a Silver Ion disc should be placed at the insertion site. PICC dressings must be changed every 7 days and as needed when a chlorhexidine-impregnated disc is in place.
 - o b) If a silver ion disc is not in place, the dressing is changed when it is no longer clean, dry, or intact.
 - o c) Tissue adhesive (for example, Secure Port IV) may be placed at the insertion site to minimize the risk of line migration.
- Assessment and documentation of insertion site, catheter and dressing integrity q 1 hour

Access frequency

- When possible, limit number of times central line is accessed to decrease risk of infection, malposition, or clotting.

Central Line placement with confirmed or suspected sepsis

- Ideal insertion of a central line occurs 48 hours after initiation of antibiotics for suspected or confirmed sepsis
- In some cases, and with attending approval, a central line may need to be placed sooner for vascular access.

Occlusion Management:

- If a central line is suspected of being completely or partially occluded:
 - o Confirm external line has no visible kinks, re-dress site if needed
 - o Consider Alteplase (TPA) treatment and use Ellucid Policy 13041: "Declotting Central Venous Catheters Using Alteplase" for additional guidelines."

Removal:

- See Ellucid Policy 13002: "Vascular Access Policy (Neonatal)" for additional central line removal instructions.
- After removal:
 - o Examine the catheter to ensure it is intact.
 - o Measure the catheter and ensure the length matches the length documented in the procedure note to ensure no catheter remains in the infant.

Complications:

Malposition:

- Upper Extremity PICC line or UVC: too deep (in Right Atrium)
 - Can lead to arrhythmias or pericardial tamponade
- UVC: too low (in hepatic veins)
 - Can cause direct injury to liver

Phlebitis

- Mechanical Phlebitis: Caused by trauma to the vein from movement of the catheter
- Treatment: Elevate Extremity, provide warmth via K-Pad for 24 hours. If no improvements, consider alternative differentials

Infection:

- Can be localized vs. systemic
- Discuss with NICU attending and/or ID attending for management guidance. Management may include obtaining a blood culture, CBC, starting antibiotics and/or removing central line

Severed Catheter

- When possible, grab the severed immediately end before it retracts into the vein/artery

- 
- Notify the NICU attending immediately

Additional Vascular Access Resources include:

- Pediatric PICC Team
- Interventional Cardiology
- Pediatric ICU Attending/Fellow
- Pediatric Surgery Attending

Medical Legal Disclaimer:

Welcome to the UC Davis Health, Department of Pediatrics, Clinical Practice Guidelines Website. All health and health-related information contained within the Site is intended chiefly for use as a resource by the Department's clinical staff and trainees in the course and scope of their approved functions/activities (although it may be accessible by others via the internet). This Site is not intended to be used as a substitute for the exercise of independent professional judgment. These clinical pathways are intended to be a guide for practitioners and may need to be adapted for each specific patient based on the practitioner's professional judgment, consideration of any unique circumstances, the needs of each patient and their family, and/or the availability of various resources at the health care institution where the patient is located. Efforts are made to ensure that the material within this Site is accurate and timely but is provided without warranty for quality or accuracy. The Regents of the University of California; University of California, Davis; University of California, Davis, Health nor any other contributing author is responsible for any errors or omissions in any information provided or the results obtained from the use of such information. Some pages within this Site, for the convenience of users, are linked to or may refer to websites not managed by UC Davis Health. UC Davis Health does not control or take responsibility for the content of these websites, and the views and opinions of the documents in this Site do not imply endorsement or credibility of the service, information or product offered through the linked sites by UC Davis Health. UC Davis Health provides limited personal permission to use the Site. This Site is limited in that you may not:

- Use, download or print material from this site for commercial use such as selling, creating course packets, or posting information on another website.
- Change or delete propriety notices from material downloaded or printed from it. • Post or transmit any unlawful, threatening, libelous, defamatory, obscene, scandalous, inflammatory, pornographic, or profane material, any propriety information belonging to others or any material that could be deemed as or encourage criminal activity, give rise to civil liability, or otherwise violate the law.
- Use the Site in a manner contrary to any applicable law.

You should assume that everything you see or read on this Site is copyrighted by University of California or others unless otherwise noted. You may download information from this Site as long as it is not used for commercial purposes, and you retain the proprietary notices. You may not use, modify, make multiple copies, or distribute or transmit the contents of this Site for public or commercial purposes without the express consent of UC Davis Health.